

**Gliklikh Yu.E.** (Voronezh, Russia) **Stochastic equations with current velocities and osmotic velocities (mean detrivatives).**

All preliminaries about mean derivatives, in particular, about current velocities (symmetric mean derivatives  $D_S$ ), osmotic velocities (antisymmetric mean derivatives  $D_A$ ) and quadratic mean derivatives  $D_2$  can be found in [1].

Let the diffusion coefficient, i.e., the field of symmetric matrices  $(a^{ij}(x))$ , be smooth, autonomous and positive definite. Since all matrices  $(\alpha^{ij}(x))$  are non-degenerate and smooth, there exist the smooth field of converse symmetric and positive definite matrices  $(\alpha_{ij})$ . Hence this field can be used as a new Riemannian metric  $\alpha(\cdot, \cdot) = \alpha_{ij} dx^i \otimes dx^j$  on  $\mathbb{R}^n$ . The volume form of metric  $\alpha(\cdot, \cdot)$  is  $\Lambda_\alpha = \sqrt{\det(\alpha_{ij}(x))} dx^1 \wedge \dots \wedge dx^n$ .

Denote by symbol  $\rho(t, x)$  the density of probablistic distribution of random element  $\xi(t)$  with respect to the volume form  $dt \wedge \Lambda_\alpha$  on  $\mathbb{R} \times \mathbb{R}^n$ , i.e., for every continuous bounded function  $f: [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$  the relation  $\int_0^T E(f(t, \xi(t))) dt = \int_0^T \left( \int_\Omega f(t, \xi(t)) d\mathbb{P} \right) dt = \int_{\mathbb{R} \times \mathbb{R}^n} f(t, x) \rho(t, x) dt \wedge \Lambda_\alpha$  holds.

Let a Borel vector field  $v(t, x)$  and a Borel field of symmetric positive semi-definite matrices  $\alpha(t, x)$  be given on  $\mathbb{R}^n$ . The system of the form

$$\begin{cases} D_S \xi(t) = v(t, \xi(t)), \\ D_2 \xi(t) = \alpha(t, \xi(t)), \end{cases} \quad (1)$$

where the equalities are fulfilled a.s., is called the first order equation with current velocities.

Everywhere below we suppose that the fields  $v$  and  $\alpha$  are smooth and all matrices  $\alpha(x)$  are autonomous and positive definite. If, in addition,  $v$ ,  $\alpha$  and partial derivatives of the coefficients of  $(a^{ij})$  satisfy the Ito inequality and the density  $\rho$  of initial value is smooth and nowhere equal to zero, it is proved that (1) has a solution (see [2]).

**Lemma 1** *Let  $\rho(t, x)$ ,  $v(t, x)$ ,  $\alpha(x)$  and  $\Lambda_\alpha$  be the same as above and the solution of (1) exists. Then the flow of vector field  $(1, v(t, x))$  on  $\mathbb{R} \times \mathbb{R}^n$  preserves the volume form  $\rho(t, x) dt \wedge \Lambda_\alpha$ , i.e., the Lie derivative  $L_{(1, v(t, x))} \rho(t, x) dt \wedge \Lambda_\alpha = 0$ .*

**Definition 1** *The system*

$$\begin{cases} D_A \xi(t) = u(t, \xi(t)) \\ D_2 \xi(t) = \alpha(\xi(t)) \end{cases}, \quad (2)$$

where  $u(t, x)$  is a Borel vector field and  $\alpha$  is as above, is called the first order differential equation with osmotic velocity.

By the use of the properties of osmotic velocity, properties of quadratic velocity and the Stokes theorem we can find  $\rho(t, x)$  that can be the density of solution of (2). Introduce  $p(t, x) = \log \rho(t, x)$ .

**Assumption 1.** *We suppose that for every  $t \in [0, T]$  the integral  $\int_{\mathbb{R} \times \mathbb{R}^n} e^{p(t, x)} dt \wedge \Lambda_\alpha$  is finite, i.e., is equal to a certain finite constant  $C(t)$  that is  $C^\infty$ -smooth in  $t$ .*

**Theorem 1** *If Assumption 1 is satisfied, under the above-mentioned hypotheses equation (2) has a solution for any initial value having the density smooth and nowhere equal to zero.*

The proof uses Lemma 1 in order to find the current velocity of solution.

1. *Gliklikh Yu.E.* Global and stochastic analysis with applications to mathematical physics.- London: Springer-Verlag.- 2011.- 460 pp.
2. *Azarina S. V., Gliklikh Yu.E.* On the Solvability of Nonautonomous Stochastic Differential Equations with Current Velocities Mathematical Notes, 2016, vol. 100, no. 1, pp. 3 – 10.